

Pure powers in recurrence sequences

KÁLMÁN LIPTAI* and TIBOR TÓMÁCS

Abstract. Let G be a linear recursive sequence of order k satisfying the recursion $G_n = A_1 G_{n-1} + \dots + A_k G_{n-k}$. In the case $k=2$ it is known that there are only finitely many perfect powers in such a sequence.

Ribenboim and McDaniel proved for sequences with $k=2$, $G_0=0$ and $G_1=1$ that in general for a term G_n there are only finitely many terms G_m such that $G_n G_m$ is a perfect square. P. Kiss proved that for any n there exists a number q_0 , depending on G and n , such that the equation $G_n G_x = w^q$ in positive integers x, w, q has no solution with $x > n$ and $q > q_0$. We show that for any n there are only finitely many $x_1, x_2, \dots, x_k, x, w, q$ positive integers such that $G_n G_{x_1} \dots G_{x_k} G_x = w^q$ and some conditions hold.

Let $R = R(A, B, R_0, R_1)$ be a second order linear recursive sequence defined by

$$R_n = AR_{n-1} + BR_{n-2} \quad (n > 1),$$

where A, B, R_0 and R_1 are fixed rational integers. In the sequel we assume that the sequence is not a degenerate one, i.e. α/β is not a root of unity, where α and β denote the roots of the polynomial $x^2 - Ax - B$.

The special cases $R(1, 1, 0, 1)$ and $R(2, 1, 0, 1)$ of the sequence R is called Fibonacci and Pell sequence, respectively.

Many results are known about relationship of the sequences R and perfect powers. For the Fibonacci sequence Cohn [2] and Wylie [23] showed that a Fibonacci number F_n is a square only when $n = 0, 1, 2$ or 12 . Pethő [12], furthermore London and Finkelstein [9,10] proved that F_n is full cube only if $n = 0, 1, 2$ or 6 . From a result of Ljunggren [8] it follows that a Pell number is a square only if $n = 0, 1$ or 7 and Pethő [12] showed that these are the only perfect powers in the Pell sequence. Similar, but more general results was showed by McDaniel and Ribenboim [11], Robbins [19,20] Cohn [3,4,5] and Pethő [15]. Shorey and Stewart [21] showed, that any non degenerate binary recurrence sequence contains only finitely many perfect powers which can be effectively determined. This results follows also from a result of Pethő [14].

* Research supported by the Hungarian National Research Science Foundation, Operating Grant Number OTKA T 16975 and 020295.

Another type of problems was studied by Ribenboim and McDaniel. For a sequence R we say that the terms R_m, R_n are in the same square-class if there exist non zero integers x, y such that

$$R_mx^2 = R_ny^2,$$

or equivalently

$$R_mR_n = t^2,$$

where t is a positive rational integer.

A square-class is called trivial if it contains only one element. Ribenboim [16] proved that in the Fibonacci sequence the square-class of a Fibonacci number F_m is trivial, if $m \neq 1, 2, 3, 6$ or 12 and for the Lucas sequence $L(1, 1, 2, 1)$ the square-class of a Lucas number L_m is trivial if $m \neq 0, 1, 3$ or 6 . For more general sequences $R(A, B, 0, 1)$, with $(A, B) = 1$, Ribenboim and McDaniel [17] obtained that each square class is finite and its elements can be effectively computed (see also Ribenboim [18]).

Further on we shall study more general recursive sequences.

Let $G = G(A_1, \dots, A_k, G_0, \dots, G_{k-1})$ be a k^{th} order linear recursive sequence of rational integers defined by

$$G_n = A_1G_{n-1} + A_2G_{n-2} + \dots + A_kG_{n-k} \quad (n > k - 1),$$

where A_1, \dots, A_k and G_0, \dots, G_{k-1} are not all zero integers. Denote by $\alpha = \alpha_1, \alpha_2, \dots, \alpha_s$ the distinct zeros of the polynomial $x^k - A_1x^{k-1} - A_2x^{k-2} - \dots - A_k$. Assume that $\alpha, \alpha_2, \dots, \alpha_s$ has multiplicity $1, m_2, \dots, m_s$ respectively and $|\alpha| > |\alpha_i|$ for $i = 2, \dots, s$. In this case, as it is known, the terms of the sequence can be written in the form

$$(1) \quad G_n = a\alpha^n + r_2(n)\alpha_2^n + \dots + r_s(n)\alpha_s^n \quad (n \geq 0),$$

where $r_i (i = 2, \dots, s)$ are polynomials of degree $m_i - 1$ and the coefficients of the polynomials and a are elements of the algebraic number field $\mathbf{Q}(\alpha, \alpha_2, \dots, \alpha_s)$. Shorey and Stewart [21] proved that the sequence G does not contain q^{th} powers if q is large enough. This result follows also from [7] and [22], where more general theorems were showed.

Kiss [6] generalized the square-class notion of Ribenboim and McDaniel. For a sequence G we say that the terms G_m and G_n are in the same q^{th} -power class if $G_mG_n = w^q$, where w, q rational integers and $q \geq 2$.

In the above mentioned paper Kiss proved that for any term G_n of the sequence G there is no terms G_m such that $m > n$ and G_n, G_m are elements of the same q^{th} -power class if q sufficiently large.

The purpose of this paper to generalize this result. We show that the under certain conditions the number of the solutions of equation

$$G_n G_{x_1} G_{x_2} \cdots G_{x_k} G_x = w^q$$

where n is fixed, are finite.

We use a well known result of Baker [1].

Lemma. *Let $\gamma_1, \dots, \gamma_v$ be non-zero algebraic numbers. Let M_1, \dots, M_v be upper bounds for the heights of $\gamma_1, \dots, \gamma_v$, respectively. We assume that M_v is at least 4. Further let b_1, \dots, b_{v-1} be rational integers with absolute values at most B and let b_v be a non-zero rational integer with absolute value at most B' . We assume that B' is at least three. Let L defined by*

$$L = b_1 \log \gamma_1 + \cdots + b_v \log \gamma_v,$$

where the logarithms are assumed to have their principal values. If $L \neq 0$, then

$$|L| > \exp(-C(\log B' \log M_v + B/B')),$$

where C is an effectively computable positive number depending on only the numbers M_1, \dots, M_{v-1} , $\gamma_1, \dots, \gamma_v$ and v (see Theorem 1 of [1] with $\delta = 1/B'$).

Theorem. *Let G be a k^{th} order linear recursive sequence satisfying the above conditions. Assume that $a \neq 0$ and $G_i \neq a\alpha^i$ for $i > n_0$. Then for any positive integer n, k and K there exists a number q_0 , depending on n, G, K and k , such that the equation*

$$(2) \quad G_n G_{x_1} G_{x_2} \cdots G_{x_k} G_x = w^q \quad (n \leq x_1 \leq \cdots \leq x_k < x)$$

in positive integer $x_1, x_2, \dots, x_k, x, w, q$ has no solution with $x_k < Kn$ and $q > q_0$.

Proof of the theorem. We can assume, without loss of generality, that the terms of the sequence G are positive. We can also suppose that $n > n_0$ and n sufficiently large since otherwise our result follows from [20] and [7].

Let $x_1, x_2, \dots, x_k, x, w, q$ positive integers satisfying (2) with the above conditions. Let ε_m be defined by

$$\varepsilon_m := \frac{1}{a} r_2(m) \left(\frac{\alpha_2}{\alpha}\right)^m + \frac{1}{a} r_3(m) \left(\frac{\alpha_3}{\alpha}\right)^m + \cdots + \frac{1}{a} r_s(m) \left(\frac{\alpha_s}{\alpha}\right)^m \quad (m \geq 0).$$

By (1) we have

$$(1 + \varepsilon_n)(1 + \varepsilon_x) \prod_{i=1}^k (1 + \varepsilon_{x_i}) a^{k+2} \alpha^{n+x+x_1+\dots+x_k} = w^q$$

from which

$$(3) \quad \begin{aligned} q \log w &= (k+2) \log a + \left(n + x + \sum_{i=1}^k x_i \right) \log \alpha + \log(1 + \varepsilon_n) \\ &+ \log(1 + \varepsilon_x) + \sum_{i=1}^k \log(1 + \varepsilon_{x_i}) \end{aligned}$$

follows. It is obvious that $x < n + x + \sum_{i=1}^k x_i < (k+2)x$. Using that $\log|1 + \varepsilon_m|$ is bounded and $\lim_{m \rightarrow \infty} \frac{1}{a} r_i(m) \left(\frac{\alpha_i}{\alpha} \right)^m = 0$ ($i = 2, \dots, s$), we have

$$(4) \quad c_1 \frac{x}{q} < \log w < c_2 \frac{x}{q}$$

where c_1 and c_2 are constants.

Let L be defined by

$$L := \left| \log \frac{w^q}{G_n G_{x_1} G_{x_2} \cdots G_{x_k} a \alpha^x} \right| = |\log(1 + \varepsilon_x)|.$$

By the definition of ε_x and the properties of logarithm function there exists a constant c_3 that

$$(5) \quad L < e^{-c_3 x}.$$

On the other hand, by the Lemma with $v = k+4$, $M_{k+4} = w$, $B' = q$ and $B = x$ we obtain the estimation

$$(6) \quad L = \left| q \log w - \log G_n - \sum_{i=1}^k \log G_{x_i} - \log a - x \log \alpha \right| > e^{-C(\log q \log w + x/q)}$$

where C depends on heights. By $x_k < Kn$ heights depend on G_n, \dots, G_{Kn} , i.e. on n, K, k and on the parameters of the recurrence. By (4), (5) and (6) we have $c_3 x < C(\log q \log w + x/q) < c_4 \log q \log w$, i.e.

$$(7) \quad x < c_5 \log q \log w$$

with some c_3, c_4, c_5 . Using (4) and (7) we get $c_6 q \log w < x < c_5 \log q \log w$, i.e. $q < c_7 \log q$, where c_6 and c_7 are constants. But this inequality does not hold if $q > q_0 = q_0(G, n, K, k)$, which proves the theorem.

References

- [1] A. BAKER, A sharpening of the bounds for linear forms in logarithms II, *Acta Arithm.* **24** (1973), 33–36.
- [2] J. H. E. COHN, On square Fibonacci numbers, *J. London Math. Soc.* **39** (1964), 537–540.
- [3] J. H. E. COHN, Squares in some recurrent sequences, *Pacific J. Math.* **41** (1972), 631–646.
- [4] J. H. E. COHN, Eight Diophantine equations, *Proc. London Math. Soc.* **16** (1966), 153–166.
- [5] J. H. E. COHN, Five Diophantine equations, *Math. Scand.* **21** (1967), 61–70.
- [6] P. KISS, Pure powers and power classes in recurrence sequences, (to appear).
- [7] P. KISS, Differences of the terms of linear recurrences, *Studia Sci. Math. Hungar.* **20** (1985), 285–293.
- [8] W. LJUNGGREN, Zur Theorie der Gleichung $x^2 + 1 = Dy^4$, *Avh. Norske Vid Akad. Oslo* **5** (1942).
- [9] J. LONDON and R. FINKELSTEIN, On Fibonacci and Lucas numbers which are perfect powers, *Fibonacci Quart.* **7** (1969) 476–481, 487, errata *ibid* **8** (1970) 248.
- [10] J. LONDON and R. FINKELSTEIN, On Mordell's equation $y^2 - k = x^3$, *Bowling Green University Press* (1973).
- [11] W. L. MCDANIEL and P. RIBENBOIM, Squares and double-squares in Lucas sequences, *C. R. Math. Acad. Sci. Soc. R. Canada* **14** (1992), 104–108.
- [12] A. PETHŐ, Full cubes in the Fibonacci sequence, *Publ. Math. Debrecen* **30** (1983), 117–127.
- [13] A. PETHŐ, The Pell sequence contains only trivial perfect powers, *Coll. Math. Soc. J. Bolyai, 60 sets, Graphs and Numbers*, Budapest, (1991), 561–568.

- [14] A. PETHŐ, Perfect powers in second order linear recurrences, *J. Number Theory* **15** (1982), 5–13.
- [15] A. PETHŐ, Perfect powers in second order recurrences, *Topics in Classical Number Theory*, Akadémiai Kiadó, Budapest, (1981), 1217–1227.
- [16] P. RIBENBOIM, Square classes of Fibonacci and Lucas numbers, *Portugaliae Math.* **46** (1989), 159–175.
- [17] P. RIBENBOIM and W. L. MCDANIEL, Square classes of Fibonacci and Lucas sequences, *Portugaliae Math.*, **48** (1991), 469–473.
- [18] P. RIBENBOIM, Square classes of $(a^n - 1)/(a - 1)$ and $a^n + 1$, *Sichuan Daxue Xunbar* **26** (1989), 196–199.
- [19] N. ROBBINS, On Fibonacci numbers of the form px^2 , where p is prime, *Fibonacci Quart.* **21** (1983), 266–271.
- [20] N. ROBBINS, On Pell numbers of the form PX^2 , where P is prime, *Fibonacci Quart.* **22** (1984), 340–348.
- [21] T. N. SHOREY and C. L. STEWART, On the Diophantine equation $ax^{2t} + bx^t y + cy^2 = d$ and pure powers in recurrence sequences, *Math. Scand.* **52** (1983), 24–36.
- [22] T. N. SHOREY and C. L. STEWART, Pure powers in recurrence sequences and some related Diophantine equations, *J. Number Theory* **27** (1987), 324–352.
- [23] O. WYLIE, In the Fibonacci series $F_1 = 1, F_2 = 1, F_{n+1} = F_n + F_{n-1}$ the first, second and twelfth terms are squares, *Amer. Math. Monthly* **71** (1964), 220–222.

KÁLMÁN LIPTAI and TIBOR TÓMACS
ESZTERHÁZY KÁROLY TEACHERS' TRAINING COLLEGE
DEPARTMENT OF MATHEMATICS
LEÁNYKA U. 4.
3301 EGER, PF. 43.
HUNGARY
E-mail: liptai@gemini.ektf.hu
tomacs@gemini.ektf.hu