

Best bounds for dispersion of ratio block sequences for certain subsets of integers

József Bukor, Peter Csiba

Department of Mathematics and Informatics
 J. Selye University, Komárno, Slovakia
bukorj@ujs.sk
csibap@ujs.sk

Submitted January 25, 2018 — Accepted May 28, 2018

Abstract

In this paper, we study the behavior of dispersion of special types of sequences which block sequence is dense.

Keywords: block sequence, dispersion, (R) -density

MSC: 11B05

1. Introduction

Denote by \mathbb{N} and \mathbb{R}^+ the set of all positive integers and positive real numbers, respectively. Let $X = \{x_1 < x_2 < x_3 < \dots\}$ be an infinite subset of \mathbb{N} . Denote by $R(X) = \{\frac{x_i}{x_j} : i, j \in \mathbb{N}\}$ the *ratio set of X* , and say that a set X is (R) -dense if $R(X)$ is (topologically) dense in the set \mathbb{R}^+ . The concept of (R) -density was introduced by T. Šalát [7].

The following sequence of finite sequences derived from X

$$\frac{x_1}{x_1}, \frac{x_1}{x_2}, \frac{x_2}{x_2}, \frac{x_1}{x_3}, \frac{x_2}{x_3}, \frac{x_3}{x_3}, \dots, \frac{x_1}{x_n}, \frac{x_2}{x_n}, \dots, \frac{x_n}{x_n}, \dots \quad (1.1)$$

is called *the block sequence* of the sequence X .

It is formed by the blocks $X_1, X_2, \dots, X_n, \dots$ where

$$X_n = \left(\frac{x_1}{x_n}, \frac{x_2}{x_n}, \dots, \frac{x_n}{x_n} \right), \quad n = 1, 2, \dots$$

is called the n -th block. This kind of block sequences was introduced by O. Strauch and J. T. Tóth [9].

For each $n \in \mathbb{N}$ consider the *step distribution function*

$$F(X_n, x) = \frac{\#\{i \leq n; \frac{x_i}{x_n} < x\}}{n},$$

and define the *set of distribution functions of the ratio block sequence*

$$G(X_n) = \left\{ \lim_{k \rightarrow \infty} F(X_{n_k}, x) \right\}.$$

The set of distribution functions of ratio block sequences was studied in [1, 2, 5, 6, 8, 12].

For every $n \in \mathbb{N}$ let

$$D(X_n) = \max \left\{ \frac{x_1}{x_n}, \frac{x_2 - x_1}{x_n}, \dots, \frac{x_{i+1} - x_i}{x_n}, \dots, \frac{x_n - x_{n-1}}{x_n} \right\},$$

the maximum distance between two consecutive terms in the n -th block. We will consider the quantity

$$\underline{D}(X) = \liminf_{n \rightarrow \infty} D(X_n),$$

(see [10]) called the *dispersion* of the block sequence (1.1) derived from X . Relations between asymptotic density and dispersion were studied in [11].

The aim of this paper is to study the behavior of dispersion of the block sequence derived from X under the assumption that $X = \bigcup_{n=1}^{\infty} (c_n, d_n) \cap \mathbb{N}$ is (R) -dense and the limit $\lim_{n \rightarrow \infty} \frac{d_n}{c_n} = s$ exists. In this case

$$\underline{D}(X) \leq \begin{cases} \frac{1}{s+1}, & \text{if } s \in \left\langle 1, \frac{1+\sqrt{5}}{2} \right\rangle, \\ \frac{1}{s^2}, & \text{if } s \in \left\langle \frac{1+\sqrt{5}}{2}, 2 \right\rangle, \\ \frac{s-1}{s^2}, & \text{if } s \in \langle 2, \infty \rangle \end{cases}$$

(see [10, Theorem 10]). This upper bound for $\underline{D}(X)$ is the best possible if $s \geq 2$ (see [4]) and in the case $\frac{1+\sqrt{5}}{2} \leq s \leq 2$ (see [3]). We prove that the above upper bound for $\underline{D}(X)$ is also optimal in the remaining case $1 \leq s < \frac{1+\sqrt{5}}{2}$, i.e. $\underline{D}(X)$ can be any number in the interval $\langle 0, \frac{1}{s+1} \rangle$.

2. Results

First, we show that there is a connection between the dispersion and the distribution functions of a ratio block sequence.

Theorem 2.1. *Let $X \subset \mathbb{N}$, and assume that the dispersion $\underline{D}(X)$ of the related block sequence is positive. Let $g \in G(X_n)$. Then g is constant on an interval of length $\underline{D}(X)$.*

Proof. Let $\varepsilon < \underline{D}(X)$ be an arbitrary positive real number. By the definition of dispersion it follows that for sufficiently large n the step distribution function $F(X_n, x)$ is constant on some interval $\left(\frac{x_i}{x_n}, \frac{x_{i+1}}{x_n}\right)$ of length $\underline{D}(X) - \varepsilon$. A simple compactness argument yields that there exist

- real numbers $\gamma, \delta \in \langle 0, 1 \rangle$ such that $\delta - \gamma \geq \underline{D}(X) - \varepsilon$,
- an increasing sequence (n_k) and a sequence (m_k) of positive integers such that $m_k < n_k$,

$$\lim_{k \rightarrow \infty} \frac{x_{m_k}}{x_{n_k}} = \gamma, \quad \lim_{k \rightarrow \infty} \frac{x_{m_k+1}}{x_{n_k}} = \delta \quad \text{and} \quad \lim_{k \rightarrow \infty} F(X_{n_k}, x) = g(x) \text{ a.e. on } \langle 0, 1 \rangle.$$

Hence g is constant on the interval (γ, δ) of length $\underline{D}(X) - \varepsilon$. Since ε can be chosen arbitrary small, and the assertion of the theorem follows. \square

The next lemma is useful for the determination of the value of the dispersion $\underline{D}(X)$ (see [10, Theorem 1]).

Lemma 2.2. *Let*

$$X = \bigcup_{n=1}^{\infty} (c_n, d_n) \cap \mathbb{N},$$

and for $n \in \mathbb{N}$ let $c_n < d_n < c_{n+1}$ be positive integers. Then

$$\underline{D}(X) = \liminf_{n \rightarrow \infty} \frac{\max\{c_{i+1} - d_i : i = 1, \dots, n\}}{d_{n+1}}.$$

For the proof of (R) -density we shall use the following lemma.

Lemma 2.3. *Denote by $(p_n), (q_n), (u_n), (v_n), (w_n)$ and (z_n) be strictly increasing sequences of positive integers satisfying*

$$p_n < q_n < u_n < v_n \quad \text{and} \quad w_n < z_n, \quad (n = 1, 2, 3, \dots).$$

Further, let

$$\left(\frac{q_n}{p_n}\right), \left(\frac{u_n}{p_n}\right), \left(\frac{v_n}{u_n}\right), \left(\frac{w_n}{u_n}\right) \text{ and } \left(\frac{z_n}{w_n}\right)$$

converge to real numbers greater than 1, moreover

$$\lim_{n \rightarrow \infty} \frac{z_n}{w_n} \geq \lim_{n \rightarrow \infty} \frac{u_n}{q_n}.$$

Then the ratio set of

$$\bigcup_n ((p_n, q_n) \cup (u_n, v_n) \cup (w_n, z_n)) \cap \mathbb{N}$$

is dense on the interval

$$\left\langle \lim_{n \rightarrow \infty} \frac{w_n}{v_n}, \lim_{n \rightarrow \infty} \frac{z_n}{p_n} \right\rangle.$$

The proof is elementary and we leave it to the reader. Let us suppose that $k \in \mathbb{N}$ is a constant. Note that the assertion of the lemma remains still true if one removes k elements from the sets $(u_n, v_n) \cap \mathbb{N}$ for all sufficiently large n .

The main result of this paper is the following.

Theorem 2.4. *Let $s \in (1, \frac{1+\sqrt{5}}{2})$ be an arbitrary real number. Then for any $\alpha \in \langle 0, \frac{1}{s+1} \rangle$ there is an (R) -dense set*

$$X = \bigcup_{n=1}^{\infty} (c_n, d_n) \cap \mathbb{N},$$

where $c_n < d_n < c_{n+1}$ are positive integers for that $\lim_{n \rightarrow \infty} \frac{d_n}{c_n} = s$ and $\underline{D}(X) = \alpha$.

Proof. It was shown in [4, Theorem 2] that the dispersion $\underline{D}(X)$ can take any number in the interval $\langle 0, \frac{s-1}{s^2} \rangle$. In what follows we suppose $\frac{s-1}{s^2} < \alpha \leq \frac{1}{s+1}$. Let us consider the function $f(x) = \frac{x-1}{sx}$. Clearly, this function is continuous and increasing on the interval $\langle 1, \infty \rangle$. Moreover

$$f(s) = \frac{s-1}{s^2} \quad \text{and} \quad f(s+1) = \frac{1}{s+1}.$$

Thus, there exists a real number $t \in (s, s+1)$ with the property

$$\frac{t-1}{st} = \alpha. \quad (2.1)$$

Write $\frac{1}{\alpha}$ in the form $s^{k+\delta}$, where k is an integer and $0 \leq \delta < 1$. The lower bound $k \geq 2$ follows from the facts that $s+1 \leq \frac{1}{\alpha}$ and $s+1 \geq s^2$ whenever $1 < s \leq \frac{1+\sqrt{5}}{2}$.

Define the set $X \subset \mathbb{N}$ by

$$X = \bigcup_{n=1}^{\infty} (A_n \cup B_n) \cap \mathbb{N},$$

where

$$A_n = \bigcup_{i=1}^k (a_{n,i}, b_{n,i}) \quad \text{and} \quad B_n = \bigcup_{j=1}^n (c_{n,j}, d_{n,j}).$$

Put $a_{1,1} = 1$ and

$$\begin{aligned} b_{n,i} &= [s.a_{n,i}] \quad \text{for } n \in \mathbb{N} \quad \text{and} \quad i = 1, 2, \dots, k, \\ a_{n,i} &= \begin{cases} d_{n-1,n-1}! & \text{for } n \geq 2, i = 1 \\ [s^\delta.b_{n,1}] + 1 & \text{for } n \in \mathbb{N}, i = 2 \\ b_{n,i-1} + 1 & \text{for } n \in \mathbb{N}, i = 3, \dots, k, \end{cases} \\ c_{n,j} &= \begin{cases} [t.b_{n,k}] + 1 & \text{for } n \in \mathbb{N}, j = 1 \\ [t.d_{n,j-1}] + 1 & \text{for } n \in \mathbb{N}, j = 2, \dots, n, \end{cases} \end{aligned}$$

$$d_{n,j} = [s.c_{n,j}] \quad \text{for } n \in \mathbb{N} \quad \text{and } j = 1, 2, \dots, n.$$

First we prove that $\underline{D}(X) = \alpha$. For sufficiently large n , by the definition of the set X we have the inequalities

$$\begin{aligned} a_{n+1,1} - d_{n,n} &> c_{n,n} - d_{n,n-1} > c_{n,n-1} - d_{n,n-2} > \dots \\ &> c_{n,3} - d_{n,2} > c_{n,2} - d_{n,1} > c_{n,1} - b_{n,k}, \end{aligned} \quad (2.2)$$

further

$$a_{n,1} - d_{n-1,n-1} < c_{n,1} - b_{n,k} \quad (2.3)$$

and

$$a_{n,2} - b_{n,1} < a_{n,1} - d_{n-1,n-1}. \quad (2.4)$$

Observe that inequality (2.3) holds if $\frac{1}{\alpha}(t-1) > 1$. In virtue of (2.1) this is equivalent with $st > 1$, which evidently holds. As

$$s^{1+\delta} - s - 1 < s^2 - s - 1$$

and $s^2 - s - 1$ is negative for $s \in (1, \frac{1+\sqrt{5}}{2})$, inequality (2.4) follows.

Now we use Lemma 1. From the inequalities (2.2-2.4) one can see that it is sufficient to study the quotients

$$\text{a) } \frac{a_{n,1} - d_{n-1,n-1}}{b_{n,k}}, \quad \text{b) } \frac{c_{n,1} - b_{n,k}}{d_{n,1}}, \quad \text{c) } \frac{c_{n,k} - d_{n,k-1}}{d_{n,k}}.$$

In case a) we see

$$\liminf_{n \rightarrow \infty} \frac{a_{n,1} - d_{n-1,n-1}}{b_{n,k}} = \liminf_{n \rightarrow \infty} \frac{a_{n,1}}{\frac{1}{\alpha}a_{n,1}} = \alpha,$$

in case b)

$$\liminf_{n \rightarrow \infty} \frac{c_{n,1} - b_{n,k}}{d_{n,1}} = \liminf_{n \rightarrow \infty} \frac{tb_{n,k} - b_{n,k}}{stb_{n,k}} = \frac{t-1}{st} = \alpha,$$

and the remaining case c) is analogous to case b).

It remains to prove that the set X is (R) -dense. Using Lemma 2 we show that the ratio set of the set X is dense on the intervals

$$\begin{aligned} \left\langle 1, \frac{1}{\alpha} \right\rangle & \text{ (for } p_n = a_{n,1}, q_n = b_{n,1}, u_n = w_n = a_{n,2}, v_n = z_n = b_{n,k}), \\ \left\langle t^i s^{i-1}, \frac{1}{\alpha} s^i t^i \right\rangle & \text{ (for } p_n = a_{n,1}, q_n = b_{n,1}, u_n = a_{n,2}, v_n = b_{n,k}, w_n = c_{n,i}, z_n = d_{n,i}). \end{aligned}$$

Hence, by $\frac{1}{\alpha} \geq s+1$ and $t < s+1$ we have

$$\left\langle 1, \frac{1}{\alpha} \right\rangle \cup \bigcup_{i=1}^{\infty} \left\langle t^i s^{i-1}, \frac{1}{\alpha} s^i t^i \right\rangle = \langle 1, \infty \rangle,$$

and therefore the (R) -density of the set X follows. □

References

- [1] BALÁŽ, V., MIŠÍK, L., STRAUCH, O., TÓTH, J. T., Distribution functions of ratio sequences, III, *Publ. Math. Debrecen*, **82** (2013), 511–529.
<https://doi.org/10.5486/pmd.2013.4770>
- [2] BALÁŽ, V., MIŠÍK, L., STRAUCH, O., TÓTH, J. T., Distribution functions of ratio sequences, IV, *Period. Math. Hung.*, **66** (2013), 1–22.
<https://doi.org/10.1007/s10998-013-4116-4>
- [3] FILIP, F., LIPTAI, K., MÁTYÁS, F., TÓTH, J. T., On the best estimations for dispersions of special ratio block sequences, *Ann. Math. Inf.*, **37** (2010), 85–93.
- [4] FILIP, F., TÓTH, J. T., On estimations of dispersions of certain dense block sequences, *Tatra Mt. Math. Publ.*, **31** (2005), 65–74.
- [5] GREKOS, G., STRAUCH, O., Distribution functions of ratio sequences, II, *Unif. Distrib. Theory* **2** (2007), 53–77.
- [6] KRĚMARSKÝ, D., MIŠÍK, L., VÁCLAVÍKOVÁ, Z., On small set of distribution functions of ratio block sequences *Unif. Distrib. Theory* **11** (2016), 165–174.
<https://doi.org/10.1515/udt-2016-0009>
- [7] ŠALÁT, T., On ratio sets of sets of natural numbers, *Acta Arith.*, **15** (1969), 273–278.
<https://doi.org/10.4064/aa-15-3-273-278>
- [8] STRAUCH, O., Distribution functions of ratio sequences. An expository paper, *Tatra Mt. Math. Publ.*, **64** (2015), 133–185.
<https://doi.org/10.1515/tmmp-2015-0047>
- [9] STRAUCH, O., TÓTH, J. T., Distribution functions of ratio sequences, *Publ. Math. Debrecen*, **58** (2001), 751–778.
- [10] TÓTH, J. T., MIŠÍK, L., FILIP, F., On some properties of dispersion of block sequences of positive integers, *Math. Slovaca*, **54** (2004), 453–464.
- [11] TÓTH, J. T., MIŠÍK, L., FILIP, F., Dispersion of ratio block sequences and asymptotic density, *Acta Arith.*, **131** (2008), 183–191.
<https://doi.org/10.4064/aa131-2-5>
- [12] TÓTH, J. T., MIŠÍK, L., FILIP, F., On ratio block sequences with extreme distribution function, *Math. Slovaca*, **59** (2009), 275–282.
<https://doi.org/10.2478/s12175-009-0123-6>