

Enumeration of Fuss-skew paths

Toufik Mansour^a, José Luis Ramírez^b

^aDepartment of Mathematics,
University of Haifa, 3498838 Haifa, Israel
tmansour@univ.haifa.ac.il

^bDepartamento de Matemáticas,
Universidad Nacional de Colombia, Bogotá, Colombia
jlr Ramirez@unal.edu.co

Abstract. In this paper, we introduce the concept of a Fuss-skew path and then we study the distribution of the semi-perimeter, area, peaks, and corners statistics. We use generating functions to obtain our main results.

Keywords: Skew Dyck path, Fuss-Catalan numbers, generating function

AMS Subject Classification: 05A15, 05A19

1. Introduction

A *skew Dyck path* is a lattice path in the first quadrant that starts at the origin, ends on the x -axis, and consists of up-steps $U = (1, 1)$, down-steps $D = (1, -1)$, and left-steps $L = (-1, -1)$, such that up and left steps do not overlap. The definition of skew Dyck path was introduced by Deutsch, Munarini, and Rinaldi [4]. Some additional results about skew Dyck path can be found in [2, 5, 8, 14].

Let s_n denote the number of skew Dyck path of semilength n , where the semilength of a path is defined as the number its up-steps. The sequence s_n is given by the combinatorial sum $s_n = \sum_{k=1}^n \binom{n-1}{k-1} c_k$, where $c_n = \frac{1}{n+1} \binom{2n}{n}$ is the n -th Catalan number. The sequence s_n appears in OEIS as A002212 [15], and its first few values are

1, 1, 3, 10, 36, 137, 543, 2219, 9285, 39587.

One way to generalize the classical Dyck paths is to regard the length of an up-step U as a parameter. Given a positive number ℓ , an ℓ -Dyck path is a lattice path in the first quadrant from $(0, 0)$ to $((\ell + 1)n, 0)$ where $n \geq 0$ using up-steps

$U_\ell = (\ell, \ell)$ and down-steps $U = (1, -1)$. For $\ell = 1$, we recover the classical Dyck path. The total number of ℓ -Dyck path with length $(\ell + 1)n$ is given by $c_\ell(n) = \frac{1}{tn+1} \binom{(t+1)n}{n}$ (cf. [1]). We will refer to ℓ -Dyck paths here as the ‘‘Fuss’’ case because the sequence $c_\ell(n)$ was first investigated by N. I. Fuss (see, for example, [7, 16] for several combinatorial interpretations for both the Catalan and Fuss-Catalan numbers).

Our focus in this paper is to introduce a Fuss analogue of the skew Dyck path. Given a positive integer ℓ , an ℓ -Fuss-skew path is a path in the first quadrant that starts at the origin, ends on the x -axis, and consists of up-steps $U_\ell = (\ell, \ell)$, down-steps $D = (1, -1)$, and left steps $L = (-1, -1)$, such that up and left steps do not overlap. Given an ℓ -Fuss-skew path P , we define the semilength of P , denote by $|P|$, as the number of up-steps of P . For example, Figure 1 shows a 3-Fuss-skew path of semilength 6. It is clear that the 1-Fuss-skew paths coincide with the skew Dyck paths. Let $\mathbb{S}_{n,\ell}$ denote the set of all ℓ -Fuss-skew path of semilength n , and $\mathbb{S}_\ell = \bigcup_{n \geq 0} \mathbb{S}_{n,\ell}$. For example, Figure 4 shows all the paths in $\mathbb{S}_{2,2}$.




Figure 1. 3-Fuss-skew path of semilength 6.

2. Counting special steps

For a given path $P \in \mathbb{S}_\ell$, we use $u(P)$, $d(P)$, and $t(P)$ to denote the number of up-steps, down-steps, and left-steps of P , respectively. In this section, we study the distribution of these parameters over \mathbb{S}_ℓ . Using these parameters, we define the generating function

$$F_\ell(x, p, q) := \sum_{P \in \mathbb{S}_\ell} x^{u(P)} p^{d(P)} q^{t(P)}.$$

For simplicity, we use F_ℓ to denote the generating function $F_\ell(x, p, q)$.

Theorem 2.1. *The generating function $F_\ell(x, p, q)$ satisfies the functional equation*

$$F_\ell = 1 + x(pF_\ell + q)^{\ell-1}(pF_\ell^2 + q(F_\ell - 1)). \tag{2.1}$$

Proof. Let \mathcal{A}_i denote the ℓ -Fuss-skew paths whose last y -coordinate is i and let A_i denote the generating function defined by

$$A_i = \sum_{P \in \mathcal{A}_i} x^{u(P)} p^{d(P)} q^{t(P)}.$$

A non-empty ℓ -Fuss-skew path can be uniquely decomposed as either $U_\ell TDP$ or $U_\ell TL$, where $U_\ell T$ is a lattice path in \mathcal{A}_1 and P is an ℓ -Fuss-skew path (see Figure 2 for a graphical representation of this decomposition). From this decomposition, we obtain the functional equation (cf. [6])

$$F_\ell = 1 + x(pA_1F_\ell + qA_1). \tag{2.2}$$




Figure 2. Decomposition of a ℓ -Fuss-skew path.

The paths of \mathcal{A}_i can be decomposed as TDP or TL , where $T \in \mathcal{A}_{i+1}$ for $i = 1, \dots, \ell - 2$ and $P \in \mathcal{S}_\ell$ (see Figure 3 for a graphical representation of this decomposition). Moreover, the paths of $\mathcal{A}_{\ell-1}$ are decomposed as P_1DP_2 or $P'L$, where $P_1, P_2, P' \in \mathcal{S}_\ell$ and P' is non-empty.




Figure 3. Decomposition of the paths in \mathcal{A}_i .

From the above decompositions, we obtain the functional equations

$$A_i = pA_{i+1}F_\ell + qA_{i+1}, \quad \text{for } i = 1, \dots, \ell - 2, \quad \text{and} \quad A_{\ell-1} = pF_\ell^2 + q(F_\ell - 1).$$

Note that in these functional equations we do not consider the first up-step because it was considered in (2.2). Therefore, we have

$$\begin{aligned} F_\ell &= 1 + x(pF_\ell + q)A_1 = 1 + x(pF_\ell + q)^2A_2 \\ &= \dots = 1 + x(pF_\ell + q)^{\ell-1}(pF_\ell^2 + q(F_\ell - 1)). \end{aligned} \quad \square$$

Let $s_\ell(n, p, q)$ denote the joint distribution over $\mathbb{S}_{n,\ell}$ for the number of down and left steps, that is,

$$s_\ell(n, p, q) = \sum_{P \in \mathbb{S}_{n,\ell}} p^{d(P)} q^{t(P)}.$$

It is clear that $F_\ell = \sum_{n \geq 0} s_\ell(n, p, q)x^n$. From the Lagrange inversion theorem (see for instance [13]), we give a combinatorial expression for the sequence $s_\ell(n, p, q)$.

Theorem 2.2. For $n \geq 1$, the sequence $s_\ell(n, p, q)$ is given by

$$\frac{1}{n} \sum_{j=0}^n \sum_{k=0}^j \binom{n}{j} \binom{j}{k} \binom{n(\ell-1)}{n-2j+k-1} p^{2n-1-2j} (2p+q)^k (p+q)^{n(\ell-2)+2j-k+1}.$$

In particular, the total number of ℓ -Fuss-skew paths of semilength n is

$$s_\ell(n) := s_\ell(n, 1, 1) = \frac{1}{n} \sum_{j=0}^n \sum_{k=0}^j \binom{n}{j} \binom{j}{k} \binom{n(\ell-1)}{n-2j+k-1} 3^k 2^{n(\ell-2)+2j-k+1}.$$

Proof. The functional equation given in Theorem 2.1 can be written as

$$Q_\ell = x(p(Q_\ell + 1) + q)^{\ell-1} (p(Q_\ell + 1)^2 + qQ_\ell),$$

where $Q_\ell = F_\ell - 1$. From the Lagrange inversion theorem, we deduce

$$\begin{aligned} [x^n]H_\ell &= \frac{1}{n} [z^{n-1}] (p(z+1) + q)^{(\ell-1)n} (p(z+1)^2 + qz)^n \\ &= \frac{1}{n} [z^{n-1}] \sum_{s \geq 0} \binom{(\ell-1)n}{s} (pz)^s (p+q)^{(\ell-1)n-s} (pz^2 + (2p+1)z + p)^n \\ &= \frac{1}{n} [z^{n-1}] \sum_{s \geq 0} \binom{(\ell-1)n}{s} (pz)^s (p+q)^{(\ell-1)n-s} \\ &\quad \times \sum_{j=0}^n \sum_{k=0}^j \binom{n}{j} \binom{j}{k} p^{n-j} ((2p+q)z)^k (pz^2)^{j-k} \\ &= \frac{1}{n} \sum_{j=0}^n \sum_{k=0}^j \binom{n}{j} \binom{j}{k} \binom{n(\ell-1)}{n-2j+k-1} p^{2n-1-2j} (2p+q)^k (p+q)^{n(\ell-2)+2j-k+1}. \quad \square \end{aligned}$$

For example, Figure 4 shows all 2-Fuss-skew paths of semilength 2 counted by the term $s_\ell(2, p, q) = 3p^4 + 6p^3q + 4p^2q^2 + pq^3$.




Figure 4. 2-Fuss-skew paths counted by $s_\ell(2, p, q)$.

From Theorem 2.2, we obtain that the total number of down-steps over the ℓ -Fuss-skew paths of semilength n is given by

$$\begin{aligned} & \left. \frac{\partial s_\ell(n, p, 1)}{\partial p} \right|_{p=1} \\ &= \frac{1}{n} \sum_{j=0}^n \sum_{k=0}^j \binom{n}{j} \binom{j}{k} \binom{n(\ell-1)}{n-2j+k-1} 2^{(\ell-2)n+2j-k} 3^{k-1} (3n(\ell+2) + k - 6j - 3). \end{aligned}$$

Moreover, the total number of left-steps over the ℓ -Fuss-skew paths of semilength n is

$$\begin{aligned} & \left. \frac{\partial s_\ell(n, 1, q)}{\partial q} \right|_{q=1} \\ &= \frac{1}{n} \sum_{j=0}^n \sum_{k=0}^j \binom{n}{j} \binom{j}{k} \binom{n(\ell-1)}{n-2j+k-1} 2^{(\ell-2)n+2j-k} 3^{k-1} (3n(\ell-2) - k + 6j + 3). \end{aligned}$$

Equation (2.1) can be explicitly solved for $\ell = 1$. In this case, we obtain the generating function

$$F_1(x, p, q) = \frac{1 - qx - \sqrt{(1 - qx)(1 - (4p + q)x)}}{2px}.$$

Moreover, the generating functions for the total number of down-steps (A026388) and left steps (A026376) over the skew-Dyck paths are respectively

$$\frac{1 - 4x + 3x^2 - \sqrt{1 - 6x + 5x^2}(1 - x)}{2x\sqrt{1 - 6x + 5x^2}}$$

and

$$\frac{1 - 3x - \sqrt{1 - 6x + 5x^2}}{2\sqrt{1 - 6x + 5x^2}}.$$

Notice that we recover some of the results of [5].

Finally, Table 1 shows the first few values of the total number of ℓ -Fuss-skew paths of semilength n .

Table 1. Values of $s_\ell(n, 1, 1)$ for $1 \leq \ell \leq 5$, $n = 1, \dots, 7$.

$\ell \setminus n$	1	2	3	4	5	6	7
$\ell = 1$	1	3	10	36	137	543	2219
$\ell = 2$	2	14	118	1114	11306	120534	1331374
$\ell = 3$	4	64	1296	29888	745856	19614464	535394560
$\ell = 4$	8	288	13568	734720	43202560	2681634816	172936069120

2.1. The width of a path

For a given path $P \in \mathbb{S}_\ell$, we define the *width* of P , denoted by $\nu(P)$, as the x -coordinate of the last point of P . For example, the width of the path given in Figure 1 is 20. We define the generating function

$$G_\ell(x, y) := G_\ell = \sum_{P \in \mathbb{S}_\ell} x^{u(P)} y^{\nu(P)}.$$

Note that each U_ℓ and D step of a path increases the width by ℓ units and 1 unit, respectively, while the left-step L decreases the width by 1 unit. Therefore, we have the functional equation

$$\begin{aligned} G_\ell &= 1 + xy^\ell (yG_\ell + y^{-1})^{\ell-1} (yG_\ell^2 + y^{-1}(G_\ell - 1)) \\ &= 1 + x(y^2G_\ell + 1)^{\ell-1} (y^2G_\ell^2 + (G_\ell - 1)). \end{aligned} \tag{2.3}$$

Let $g_\ell(n, y)$ denote the distribution over $\mathbb{S}_{n,\ell}$ for the width parameter, i.e.,

$$g_\ell(n, y) = \sum_{P \in \mathbb{S}_{n,\ell}} y^{\nu(P)}.$$

From the functional equation (2.3) and the Lagrange inversion theorem, we obtain the following theorem.

Theorem 2.3. For $n \geq 1$, the sequence $g_\ell(n, y)$ is given by

$$\frac{1}{n} \sum_{j=0}^n \sum_{k=0}^j \binom{n}{j} \binom{j}{k} \binom{n(\ell-1)}{n-2j+k-1} y^{4(n-j)-2} (y^2+1)^{n(\ell-2)+2j-k+1} (2y^2+1)^k.$$

For example, $g_2(2, y) = y^2 + 4y^4 + 6y^6 + 3y^8$. This polynomial can be found from the paths in Figure 4. For $\ell = 1$, we obtain the explicit generating function with respect to the width of a skew Dyck path.

$$G_1(x, y) = \frac{1 - x - \sqrt{(1-x)(1-x-4xy^2)}}{2xy^2}.$$

3. Number of peaks

For a given path $P \in \mathbb{S}_\ell$, we define the *peaks* of P , denoted by $\rho(P)$, as the number of subpaths of the form $U_\ell D$ (for counting peaks in a Dyck path, for example, see [9, 11]). For example, the number of peaks of the path given in Figure 1 is 5. We define the generating function

$$P_\ell(x, y) := P_\ell = \sum_{P \in \mathbb{S}_\ell} x^{u(P)} y^{\rho(P)}.$$

Theorem 3.1. *The generating function $P_\ell(x, y)$ satisfies the functional equation*

$$P_\ell = 1 + x(P_\ell + 1)^{\ell-1}((P_\ell - 1 + y)P_\ell + (P_\ell - 1)).$$

Proof. Let C_i denote the generating function defined by $C_i = \sum_{P \in \mathcal{A}_i} x^{u(P)} y^{\rho(P)}$. From the decomposition given for the ℓ -Fuss-skew paths, we have the equation $P_\ell = 1 + x(C_1 P_\ell + C_1)$. Moreover,

$$\begin{aligned} C_i &= C_{i+1} P_\ell + C_{i+1}, \quad \text{for } i = 1, \dots, \ell - 2, \text{ and} \\ C_{\ell-1} &= (P_\ell - 1 + y)P_\ell + (P_\ell - 1). \end{aligned}$$

From these relations, we obtain the desired result. □

Let $p_\ell(n, y)$ denote the distribution over \mathbb{S}_n for the peaks statistic, i.e.,

$$p_\ell(n, y) = \sum_{P \in \mathbb{S}_n} y^{\rho(P)}.$$

From the Lagrange inversion theorem, we deduce the following result.

Theorem 3.2. *For $n \geq 1$, we have*

$$p_\ell(n, y) = \frac{1}{n} \sum_{j=0}^n \sum_{k=0}^j \binom{n}{j} \binom{j}{k} \binom{n(\ell-1)}{n-2j+k-1} 2^{n(\ell-2)+2j-k+1} y^{n-j} (y+2)^k.$$

In particular, the total number of peaks in all ℓ -Fuss-skew paths of semilength n is

$$\begin{aligned} & \left. \frac{\partial p_\ell(n, y)}{\partial y} \right|_{y=1} \\ &= \frac{1}{n} \sum_{j=0}^n \sum_{k=0}^j \binom{n}{j} \binom{j}{k} \binom{n(\ell-1)}{n-2j+k-1} 2^{n(\ell-2)+2j-k+1} 3^{k-1} (3(n-j) + k). \end{aligned}$$

For example, $p_2(2, y) = 8y + 6y^2$. This polynomial can be found from the paths in Figure 4. For $\ell = 1$ we obtain the generating function

$$P_1(x, y) = \frac{1 - xy - \sqrt{(1 - xy)^2 - 4(1 - x)x}}{2x}.$$

Moreover, the generating function for the total number of peaks is

$$\frac{1 - x - \sqrt{1 - 6x + 5x^2}}{2\sqrt{1 - 6x + 5x^2}}.$$

Table 2 shows the first few values of the number of peaks in ℓ -Fuss-skew paths of semilength n .

Table 2. Total number of peaks in \mathbb{S}_ℓ .

$\ell \setminus n$	1	2	3	4	5	6	7
$\ell = 1$	1	4	17	75	339	1558	7247
$\ell = 2$	2	20	226	2696	33138	415164	5270850
$\ell = 3$	4	96	2672	78848	2400896	74568704	2347934464
$\ell = 4$	8	448	29440	2054144	147986432	10878189568	810813030400

4. Number of corners

For a given path $P \in \mathbb{S}_\ell$, we define a *corner* of P as a right angle caused by two consecutive steps in the graph of P . For example, the path given in Figure 5 has 4 corners, depicted in red. This statistic has been studied in other combinatorial structures as integer partitions [3], compositions [10], and bargraphs [12].




Figure 5. Corners of a path.

Let $\tau(P)$ denote the number of corners of P . We define the bivariate generating function

$$W_\ell(x, y) := W_\ell = \sum_{P \in \mathbb{S}_\ell} x^{u(P)} y^{\tau(P)}.$$

In this section, we analyze the cases $\ell = 1$ and $\ell = 2$. We leave as an open question the case $\ell \geq 3$.

Theorem 4.1. *The generating function $W_1(x, y)$ satisfies the functional equation*

$$xy(1 + y)W_1^3 - (2 - x(2 - y^2))W_1^2 + 3(1 - x)W_1 + x - 1 = 0.$$

Proof. Let \mathcal{D} and \mathcal{L} denote the skew Dyck paths whose last step is a down-step or a left-step, respectively. Let D and L denote the generating functions defined by

$$D = \sum_{P \in \mathcal{D}} x^{u(P)} y^{\tau(P)} \quad \text{and} \quad L = \sum_{P \in \mathcal{L}} x^{u(P)} y^{\tau(P)}.$$

A non-empty skew Dyck path can be uniquely decomposed as either UT_1L or UT_2DT_3 , where T_1, T_2 , and T_3 are lattice paths in \mathbb{S}_1 with T_1 non-empty. In the first case, T_1 has two options: the last step is a down-step or a left step, see Figure 6. Then, this case contributes to the generating function the term $x(yD + L)$.




Figure 6. Decomposition of a skew Dyck path.

On the other hand, T_2 can be an empty path or a path in \mathcal{D} or \mathcal{L} . If T_3 is empty, then this case contributes to the generating function the term $x(y + D + Ly)$. On the other hand, if the path T_3 is non-empty, then this case contributes to the generating function the term $x(y + D + yL)y(W_1 - 1)$, see Figure 7. Summarizing these cases, we obtain the functional equation

$$W_1 = 1 + x(yD + L) + x(y + D + yL)(1 + y(W_1 - 1)).$$

From a similar argument, we obtain the equations

$$D = x(y + D + yL)(1 + yD) \quad \text{and} \quad L = x(yD + L) + x(y + D + yL)(yL).$$




Figure 7. Decomposition of a skew Dyck path.

Using the Gröbner basis on the polynomial equations for W_1, D , and L , we obtain the desired result. \square

We can use a symbolic software computation to obtain the first few terms of the formal power series of $W_1(x, y)$ as follows:

$$W_1(x, y) = 1 + xy + x^2(y + y^2 + y^3) + x^3(y + 2y^2 + 4y^3 + 2y^4 + y^5) + x^4(y + 3y^2 + 9y^3 + 9y^4 + 10y^5 + 3y^6 + y^7) + \dots$$

From the equation given in Theorem 4.1, we obtain

$$3xS^3(x) + 6xS^2(x)K(x) - 2xS^2(x) - 2(2 - x)S(x)K(x) + 3(1 - x)K(x) = 0,$$

where $K(x)$ is the generating function for the total number of corners in skew Dyck paths and $S(x) = (1 - x - \sqrt{1 - 6x + 5x^2})/(2x)$ is the generating function for the number of the skew Dyck paths. Solving the above equation, we obtain the generating function

$$K(x) = \frac{2(1 - x)(3 + x)x}{(1 - x)(3 - 2x)(1 - 5x) + (3 - 11x + 4x^2)\sqrt{1 - 6x + 5x^2}} = x + 6x^2 + 30x^3 + 145x^4 + 695x^5 + 3327x^6 + 15945x^7 + \dots$$

Theorem 4.2. *The generating function $W_2(x, y)$ satisfies the functional equation*

$$\begin{aligned} & x^2y^4(1 + y)^3W_2^6 - xy^2(1 + y)^2(1 - x(1 + 6y + y^2 - 3y^3))W_2^5 \\ & + xy(-4 - 7y + 3y^3 + x(1 + y)^2(4 + 9y - 11y^2 - 6y^3 + 3y^4))W_2^4 \\ & + (4 - 2x(1 + y)^2(4 - 7y + y^2) - x^2(1 + y)^2(-4 + 2y + 21y^2 - 8y^3 - 5y^4 + y^5))W_2^3 \\ & + (-12 - x^2(1 + y)^2(8 + 4y - 18y^2 + 4y^3 + y^4) - 2x(-10 - 9y + 6y^2 + 6y^3 + y^4))W_2^2 \\ & + (12 + x^2(1 + y)^2(5 + 4y - 7y^2 + y^3) + x(-17 - 16y + 2y^2 + 4y^3 + 3y^4))W_2 \\ & + (-4 + x^2(1 + y)^2(-1 - y + y^2) + x(5 + 4y - y^4)) = 0. \end{aligned}$$

Proof. Let \mathcal{D}_2 and \mathcal{L}_2 denote the 2-Fuss-skew paths whose last step is a down-step or a left-step, respectively. Let D_2 and L_2 denote the generating functions defined by

$$D_2 = \sum_{P \in \mathcal{D}_2} x^{u(P)}y^{\tau(P)} \quad \text{and} \quad L_2 = \sum_{P \in \mathcal{L}_2} x^{u(P)}y^{\tau(P)}.$$

From a similar argument as in the proof of Theorem 4.1, we obtain the system of polynomial equations

$$\begin{aligned} W_2 &= 1 + x((y + yD_2 + L_2)(1 + y^2D_2 + yL_2)(1 + y(W_2 - 1)) + (D_2 + yL_2) \\ & \quad + (D_2 + yL_2)y(1 + y(W_2 - 1)) + (y + yD_2 + L_2)(y + yD_2 + y^2L_2)), \\ D_2 &= x((y + yD_2 + L_2)(1 + y^2D_2 + yL_2)yD_2 + (D_2 + yL_2) + (D_2 + yL_2)y(yD_2) \\ & \quad + (y + yD_2 + L_2)(y + yD_2 + y^2L_2)), \\ L_2 &= x((y + yD_2 + L_2)(1 + y^2D_2 + yL_2)(1 + yL_2) + (D_2 + yL_2)y(1 + yL_2)). \end{aligned}$$

By using the Gröbner basis, we obtain the desired result. \square

Expanding with *Mathematica* the functional equation for W_2 , we find

$$W_2(x, y) = 1 + (y + y^2)x + (y + 3y^2 + 5y^3 + 4y^4 + y^5)x^2 + (y + 5y^2 + 16y^3 + 27y^4 + 33y^5 + 25y^6 + 9y^7 + 2y^8)x^3 + \dots .$$

Moreover, the first few terms of the total number of corners in \mathbb{S}_2 are

$$3x + 43x^2 + 561x^3 + 7209x^4 + 92703x^5 + 1197151x^6 + 15532917x^7 + 202428373x^8 + \dots .$$

From Figure 4 one can verify that there are 43 corners over all paths in $\mathbb{S}_{2,2}$.

5. Other generalization

Let \mathbb{H}_ℓ denote the skew Dyck paths where left steps are below the line $y = \ell$. In particular, \mathbb{H}_0 are the Dyck path and \mathbb{H}_∞ are the skew Dyck path. We define the generating function

$$H_\ell(x, p, q) := \sum_{P \in \mathbb{H}_\ell} x^{u(P)} p^{d(P)} q^{t(P)} .$$

For simplicity, we use H_ℓ to denote the generating function $H_\ell(x, p, q)$.

Theorem 5.1. *For $\ell \geq 1$, we have*

$$H_\ell = 1 + qx(H_{\ell-1} - 1) + pxH_{\ell-1}H_\ell, \tag{5.1}$$

with the initial value $H_0 = \frac{1 - \sqrt{1 - 4px}}{2px}$.

Proof. A non-empty skew Dyck path in \mathbb{H}_ℓ can be decomposed as UT_1L or UT_2DT_3 , where $T_1, T_2 \in \mathbb{H}_{\ell-1}$ with T_1 a non-empty path, and $T_3 \in \mathbb{H}_\ell$. From this decomposition follows the functional equation. \square

Recall that the m th Chebyshev polynomial of the second kind satisfies the recurrence relation $U_m(t) = 2tU_{m-1}(t) - U_{m-2}(t)$ with $U_0(t) = 1$ and $U_1(t) = 2t$. Thus by induction on ℓ and Theorem 5.1, we obtain the following result.

Theorem 5.2. *Let $t = \frac{1+qx}{2\sqrt{x(p+q-pqx)}}$ and $r = \sqrt{x(p+q-pqx)}$. The generating function H_ℓ is given by*

$$\frac{(qxU_{n-1}(t) - rU_{n-2}(t))C(px) + (1 - qx)U_{n-1}(t)}{U_{n-1}(t) - rU_{n-2}(t) - pxU_{n-1}(t)C(px)},$$

where U_m is the m th Chebyshev polynomial of the second kind and $C(x) = \frac{1 - \sqrt{1 - 4x}}{2x}$ the generating function for the Catalan numbers $\frac{1}{n+1} \binom{2n}{n}$.

The generating functions for the total number of skew Dyck path in \mathbb{H}_ℓ for $\ell = 1, 2, 3$ are

$$H_1(x, 1, 1) = \frac{3 - 2x - \sqrt{1 - 4x}}{1 + \sqrt{1 - 4x}},$$

$$H_2(x, 1, 1) = \frac{1 + 2x - 2x^2 - (1 - 2x)\sqrt{1 - 4x}}{1 - x - 2(1 - x)x + (1 + x)\sqrt{1 - 4x}},$$

$$H_3(x, 1, 1) = \frac{1 - 3x + 7x^2 - 4x^3 + (1 + x - 3x^2)\sqrt{1 - 4x}}{1 - 4x + 2x^3 + (1 + 2x^2)\sqrt{1 - 4x}}.$$

Acknowledgements. We thanks Mark Shattuck for his comments on the previous version of the paper.

References

- [1] J.-C. AVAL: *Multivariate Fuss-Catalan numbers*, Discrete Math. 308 (2008), pp. 4660–4669, DOI: <https://doi.org/10.1016/j.disc.2007.08.100>.
- [2] J. L. BARIL, J. L. RAMÍREZ, L. M. SIMBAQUEBA: *Counting prefixes of skew Dyck paths*, J. Integer Seq. Article 21.8.2 (2021).
- [3] A. BLECHER, C. BRENNAN, A. KNOPFMACHER, T. MANSOUR: *Counting corners in partitions*, Ramanujan J. 39 (2016), pp. 201–224, DOI: <https://doi.org/10.1007/s11139-014-9666-4>.
- [4] E. DEUTSCH, E. MUNARINI, S. RINALDI: *Skew Dyck paths*, J. Statist. Plann. Inference 140 (2010), pp. 2191–2203, DOI: <https://doi.org/10.1016/j.jspi.2010.01.015>.
- [5] E. DEUTSCH, E. MUNARINI, S. RINALDI: *Skew Dyck paths, area, and superdiagonal bargraphs*, J. Statist. Plann. Inference 140 (2010), pp. 1550–1526, DOI: <https://doi.org/10.1016/j.jspi.2009.12.013>.
- [6] P. FLAJOLET, R. SEDGEWICK: *Analytic Combinatorics*, Cambridge University Press, 2009.
- [7] S. HEUBACH, N. Y. LI, T. MANSOUR: *Staircase tilings and k-Catalan structures*, Discrete Math. 308 (2008), pp. 5954–5964, DOI: <https://doi.org/10.1016/j.disc.2007.11.012>.
- [8] Q. L. LU: *Skew Motzkin paths*, Acta Math. Sin. (Engl. Ser.) 33 (2017), pp. 657–667, DOI: <https://doi.org/10.1007/s10114-016-5292-y>.
- [9] T. MANSOUR: *Counting peaks at height k in a Dyck path*, J. Integer Seq. Article 02.1.1 (2002).
- [10] T. MANSOUR, A. S. SHABANI, M. SHATTUCK: *Counting corners in compositions and set partitions presented as bargraphs*, J. Difference Equ. Appl. 24 (2018), pp. 992–1015, DOI: <https://doi.org/10.1080/10236198.2018.1444760>.
- [11] T. MANSOUR, M. SHATTUCK: *Counting Dyck paths according to the maximum distance between peaks and valleys*, J. Integer Seq. Article 12.1.1 (2012).
- [12] T. MANSOUR, G. YILDIRIM: *Enumerations of bargraphs with respect to corner statistics*, Appl. Anal. Discrete Math. 14 (2020), pp. 221–238, DOI: <https://dx.doi.org/10.2298/AADM181101009M>.
- [13] D. MERLINI, R. SPRUGNOLI, M. C. VERRI: *Lagrange inversion: when and how*, Acta Appl. Math. 94 (2006), pp. 233–249, DOI: <https://doi.org/10.1007/s10440-006-9077-7>.
- [14] H. PRODINGER: *Partial skew Dyck paths - A kernel method approach*, arXiv:2108.09785v1 (2021), DOI: <https://arxiv.org/abs/2108.09785v2>.
- [15] N. J. A. SLOANE, The On-Line Encyclopedia of Integer Sequences, DOI: <https://oeis.org/>.
- [16] R. STANLEY: *Catalan Numbers*, Cambridge University Press, 2015.