

A note on the exponential Diophantine equation $(a^x - 1)(b^y - 1) = az^2$

Yasutsugu Fujita^a, Maohua Le^b

^aDepartment of Mathematics, College of Industrial Technology, Nihon University,
2-11-1 Shin-ei, Narashino, Chiba, Japan
fujita.yasutsugu@nihon-u.ac.jp

^bInstitute of Mathematics, Lingnan Normal College,
Zhanjiang, Guangdong, 524048 China
lemaohua2008@163.com

Abstract. Let a, b be fixed positive integers such that $(a \bmod 8, b \bmod 8) \in \{(0, 3), (0, 5), (2, 3), (2, 5), (4, 3), (6, 5)\}$. In this paper, using elementary methods with some classical results for Diophantine equations, we prove the following three results: (i) The equation $(*)$ $(a^x - 1)(b^y - 1) = az^2$ has no positive integer solutions (x, y, z) with $2 \nmid x$ and $x > 1$. (ii) If $a = 2$ and $b \equiv 5 \pmod{8}$, then $(*)$ has no positive integer solutions (x, y, z) with $2 \nmid x$. (iii) If $a = 2$ and $b \equiv 3 \pmod{8}$, then the positive integer solutions (x, y, z) of $(*)$ with $2 \nmid x$ are determined. These results improve the recent results of R.-Z. Tong: On the Diophantine equation $(2^x - 1)(p^y - 1) = 2z^2$, Czech. Math. J. 71 (2021), 689–696. Moreover, under the assumption that a is a square, we prove that $(*)$ has no positive integer solutions (x, y, z) even with $2 \mid x$ in some cases.

Keywords: polynomial-exponential Diophantine equation, Pell's equation, generalized Ramanujan-Nagell equation

AMS Subject Classification: 11D61

1. Introduction

Let \mathbb{N} be the set of all positive integers. Let a, b be fixed positive integers with $\min\{a, b\} > 1$. In 2000, L. Szalay [7] completely solved the equation

$$(2^x - 1)(3^x - 1) = z^2, \quad x, z \in \mathbb{N}. \quad (1.1)$$

He proved that (1.1) has no solutions (x, z) . Since then, this result has led to a series of related studies for the equation

$$(a^x - 1)(b^x - 1) = z^2, \quad x, z \in \mathbb{N} \quad (1.2)$$

(see [3]). Obviously, the solution of (1.2) involves a system of generalized Ramanujan-Nagell equations. Recently, R.-Z. Tong [8] discussed the equation

$$(2^x - 1)(p^y - 1) = 2z^2, \quad x, y, z \in \mathbb{N}, \quad (1.3)$$

where p is an odd prime with $p \equiv \pm 3 \pmod{8}$. He proved the following two results: (i) (1.3) has no solutions (x, y, z) with $2 \nmid x$, $2 \mid y$ and $y > 4$. (ii) If $p \neq 2g^2 + 1$, where g is an odd positive integer, then (1.3) has no solutions (x, y, z) with $2 \nmid x$. In this paper, we will discuss the generalized form of (1.3) as follows:

$$(a^x - 1)(b^y - 1) = az^2, \quad x, y, z \in \mathbb{N}. \quad (1.4)$$

For any positive integer n , let r_n, s_n be the positive integers satisfying

$$r_n + s_n\sqrt{2} = (3 + 2\sqrt{2})^n. \quad (1.5)$$

For any odd positive integer m , let R_m, S_m be the positive integers satisfying

$$R_m + S_m\sqrt{2} = (1 + \sqrt{2})^m. \quad (1.6)$$

Using elementary methods with some classical results for Diophantine equations, we prove the following results:

Theorem 1.1. *If*

$$(a \pmod{8}, b \pmod{8}) \in \{(0, 3), (0, 5), (2, 3), (2, 5), (4, 3), (6, 5)\}, \quad (1.7)$$

then (1.4) has no solutions (x, y, z) with $2 \nmid x$ and $x > 1$.

Theorem 1.2. *If $a = 2$ and $b \equiv 5 \pmod{8}$, then (1.4) has no solutions (x, y, z) with $2 \nmid x$. If $a = 2$ and $b \equiv 3 \pmod{8}$, then (1.4) has only the following solutions (x, y, z) with $2 \nmid x$:*

(i) $b = 3$, $(x, y, z) = (1, 1, 1)$, $(1, 2, 2)$ and $(1, 5, 11)$.

(ii) $b = 2g^2 + 1$, $(x, y, z) = (1, 1, g)$, where g is an odd positive integer with $g > 1$.

(iii) $b = r_m$, $(x, y, z) = (1, 2, s_m)$, where m is an odd positive integer with $m > 1$.

Theorem 1.3. *Let $N(a, b)$ denote the number of solutions (x, y, z) of (1.4) with $2 \nmid x$. If $a = 2$ and $b \equiv 3 \pmod{8}$, then*

$$N(2, b) = \begin{cases} 3, & \text{if } b = 3, \\ 2, & \text{if } b = 2g^2 + 1 \text{ and } g = R_m \text{ with } m > 1, \\ 1, & \text{if } b = 2g^2 + 1 \text{ and } g \neq R_m, \\ 0, & \text{otherwise.} \end{cases}$$

Obviously, the above theorems improve the result of [8].

The following results concern the solvability of (1.4) including even the case where $2 \mid x$.

Theorem 1.4. *If $(a \bmod 8, b \bmod 8) \in \{(0, 3), (0, 5), (4, 3)\}$ and a is a square, then (1.4) has no solutions (x, y, z) with $x > 1$.*

Theorem 1.5. *Assume that one of the following conditions holds:*

(i) $a = 4$ and either $b = 3$ or b has a prime divisor p with $p \equiv 11 \pmod{24}$.

(ii) $a = 16$ and either $b \in \{3, 5\}$ or b has a prime divisor p with

$$p \equiv 11, 13, 29, 37, 43, 59, 67 \text{ or } 101 \pmod{120}.$$

Then, (1.4) has no solutions.

2. Preliminaries

Let D be a nonsquare positive integer, and let D_1, D_2 be positive integers such that $D_1 > 1$, $D_1 D_2 = D$ and $\gcd(D_1, D_2) = 1$. By the basic properties of Pell's equation (see [5, 10] and [4, Lemma 1]), we obtain the following two lemmas immediately.

Lemma 2.1. *The equation*

$$u^2 - Dv^2 = 1, \quad u, v \in \mathbb{N} \tag{2.1}$$

has solutions (u, v) , and it has a unique solution (u_1, v_1) such that $u_1 + v_1\sqrt{D} \leq u + v\sqrt{D}$, where (u, v) runs through all solutions of (2.1). The solution (u_1, v_1) is called the least solution of (2.1). For any positive integer n , let $u_n + v_n\sqrt{D} = (u_1 + v_1\sqrt{D})^n$. Then we have

(i) $(u, v) = (u_n, v_n)$ ($n = 1, 2, \dots$) are all solutions of (2.1).

(ii) If $2 \mid n$, then each prime divisor p of u_n satisfies $p \equiv \pm 1 \pmod{8}$.

(iii) If $2 \nmid n$, then $u_1 \mid u_n$.

Lemma 2.2. *If the equation*

$$D_1 U^2 - D_2 V^2 = 1, \quad U, V \in \mathbb{N} \tag{2.2}$$

has solutions (U, V) , then it has a unique solution (U_1, V_1) such that $U_1\sqrt{D_1} + V_1\sqrt{D_2} \leq U\sqrt{D_1} + V\sqrt{D_2}$, where (U, V) runs through all solutions of (2.2). The solution (U_1, V_1) is called the least solution of (2.2). For any odd positive integer m , let $U_m\sqrt{D_1} + V_m\sqrt{D_2} = (U_1\sqrt{D_1} + V_1\sqrt{D_2})^m$. Then we have

(i) $(U, V) = (U_m, V_m)$ ($m = 1, 3, \dots$) are all solutions of (2.2).

(ii) $u_1 + v_1\sqrt{D} = (U_1\sqrt{D_1} + V_1\sqrt{D_2})^2$, where (u_1, v_1) is the least solution of (2.1).

For any positive integer l , let $\text{ord}_2(l)$ denote the order of 2 in the factorization of l .

Lemma 2.3. *If (2.2) has solutions (U, V) , then every solution (U, V) of (2.2) satisfies $\text{ord}_2(D_1U^2) = \text{ord}_2(D_1U_1^2)$, where (U_1, V_1) is the least solution of (2.2).*

Proof. By (i) of Lemma 2.2, there exists an odd positive integer m which makes $U\sqrt{D_1} + V\sqrt{D_2} = (U_1\sqrt{D_1} + V_1\sqrt{D_2})^m$, whence we get

$$U = U_1 \sum_{i=0}^{(m-1)/2} \binom{m}{2i} (D_1U_1^2)^{(m-1)/2-i} (D_2V_1^2)^i. \quad (2.3)$$

Since $D_1U_1^2 - D_2V_1^2 = 1$ implies that $D_1U_1^2$ and $D_2V_1^2$ have opposite parity, we have

$$2 \nmid \sum_{i=0}^{(m-1)/2} \binom{m}{2i} (D_1U_1^2)^{(m-1)/2-i} (D_2V_1^2)^i. \quad (2.4)$$

Hence, by (2.3) and (2.4), we get $\text{ord}_2(U) = \text{ord}_2(U_1)$. It implies that $\text{ord}_2(D_1U^2) = \text{ord}_2(D_1U_1^2)$. The lemma is proved. \square

Lemma 2.4. *Let r_n, s_n be defined as in (1.5). Then $(u, v) = (r_n, s_n)$ ($n = 1, 2, \dots$) are all solutions of the equation*

$$u^2 - 2v^2 = 1, \quad u, v \in \mathbb{N}, \quad (2.5)$$

and

$$r_n \equiv \begin{cases} 1 \pmod{8}, & \text{if } 2 \mid n, \\ 3 \pmod{8}, & \text{if } 2 \nmid n. \end{cases} \quad (2.6)$$

Proof. Since $(u_1, v_1) = (3, 2)$ is the least solution of (2.5), by (i) of Lemma 2.1, we see from (1.5) that $(u, v) = (r_n, s_n)$ ($n = 1, 2, \dots$) are all solutions of (2.5). By (1.5) we have

$$r_n = \sum_{i=0}^{[n/2]} \binom{m}{2i} 3^{n-2i} \cdot 8^i,$$

where $[n/2]$ is the integer part of $n/2$. It follows that

$$r_n \equiv 3^n \pmod{8},$$

whence we obtain (2.6). The lemma is proved. \square

Lemma 2.5. *For any odd positive integer m , we have $r_m = 2R_m^2 + 1$, where r_m, R_m are defined as in (1.5) and (1.6) respectively.*

Proof. Since $3 + 2\sqrt{2} = (1 + \sqrt{2})^2$ and $3 - 2\sqrt{2} = (1 - \sqrt{2})^2$, by (1.5) and (1.6), we have

$$\begin{aligned} r_m &= \frac{1}{2} \left((3 + 2\sqrt{2})^m + (3 - 2\sqrt{2})^m \right) = \frac{1}{2} \left((1 + \sqrt{2})^{2m} + (1 - \sqrt{2})^{2m} \right) \\ &= \frac{1}{2} \left(\left((1 + \sqrt{2})^m + (1 - \sqrt{2})^m \right)^2 - 2(1 + \sqrt{2})^m (1 - \sqrt{2})^m \right) \\ &= \frac{1}{2} \left((2R_m)^2 + 2 \right) = 2R_m^2 + 1. \end{aligned}$$

The lemma is proved. □

Lemma 2.6 ([9]). *The equation*

$$2X^2 + 1 = Y^3, \quad X, Y \in \mathbb{N}$$

has no solutions (X, Y) .

Lemma 2.7 ([6]). *The equation*

$$2X^2 + 1 = Y^q, \quad X, Y \in \mathbb{N}, \quad q \text{ is an odd prime with } q > 3$$

has only the solution $(X, Y, q) = (11, 3, 5)$.

Lemma 2.8 ([1, 2]). *The equation*

$$X^4 - DY^2 = 1, \quad X, Y \in \mathbb{N}$$

has solutions (X, Y) if and only if either $X^2 = u_1$ or $X^2 = 2u_1^2 - 1$.

Lemma 2.9. *The equation*

$$2X^2 + 1 = Y^t, \quad X, Y, t \in \mathbb{N}, \quad t > 2 \tag{2.7}$$

has only the solution $(X, Y, t) = (11, 3, 5)$.

Proof. Let (X, Y, t) be a solution of (2.7), and let q be the largest prime divisor of t . By Lemmas 2.6 and 2.7, (2.7) has only the solution $(X, Y, t) = (11, 3, 5)$ with $q \geq 3$. Since $t > 2$, if $q = 2$, then $4 \mid t$ and the equation

$$(X')^4 - 2(Y')^2 = 1, \quad X', Y' \in \mathbb{N} \tag{2.8}$$

has a solution $(X', Y') = (Y'^{t/4}, X)$. However, since the least solution of (2.5) is $(u_1, v_1) = (3, 2)$, neither $u_1 = 3$ nor $2u_1^2 - 1 = 17$ is a square. By Lemma 2.8, (2.8) has no solutions (X', Y') . Therefore, (2.7) has no solutions (X, Y, t) with $q = 2$. The lemma is proved. □

3. Proof of Theorem 1.1

In this section, we assume that (1.7) holds and that (x, y, z) is a solution of (1.4) with $2 \nmid x$ and $x > 1$. Then we have

$$x \geq 3. \quad (3.1)$$

Since $\gcd(a, a^x - 1) = 1$, by (1.4), we get

$$a^x - 1 = df^2, \quad b^y - 1 = adg^2, \quad z = dfg, \quad d, f, g \in \mathbb{N}. \quad (3.2)$$

By the first equality of (3.2), we have

$$\gcd(a, d) = 1. \quad (3.3)$$

Since $2 \mid a$, by (3.1) and the first equality of (3.2), we get $2 \nmid f$ and

$$d \equiv df^2 \equiv a^x - 1 \equiv 0 - 1 \equiv 7 \pmod{8}. \quad (3.4)$$

Hence, we see from (3.4) that

$$d \text{ is not a square.} \quad (3.5)$$

On the other hand, substituting (3.4) into the second equality of (3.2), we have

$$b^y \equiv 1 + 7ag^2 \equiv \begin{cases} 1 \pmod{8}, & \text{if } a \equiv 0 \pmod{8} \text{ or } 2 \mid g, \\ 7 \pmod{8}, & \text{if } a \equiv 2 \pmod{8} \text{ and } 2 \nmid g, \\ 5 \pmod{8}, & \text{if } a \equiv 4 \pmod{8} \text{ and } 2 \nmid g, \\ 3 \pmod{8}, & \text{if } a \equiv 6 \pmod{8} \text{ and } 2 \nmid g. \end{cases} \quad (3.6)$$

Further, since $b \equiv \pm 3 \pmod{8}$, we get

$$b^y \equiv \begin{cases} 1 \pmod{8}, & \text{if } 2 \mid y, \\ \pm 3 \pmod{8}, & \text{if } 2 \nmid y. \end{cases} \quad (3.7)$$

Therefore, in view of (1.7), comparing (3.6) and (3.7), we obtain

$$2 \mid y. \quad (3.8)$$

We see from (3.8) and the second equality of (3.2) that the equation

$$u^2 - adv^2 = 1, \quad u, v \in \mathbb{N} \quad (3.9)$$

has a solution

$$(u, v) = (b^{y/2}, g). \quad (3.10)$$

By (3.3) and (3.5), ad is a nonsquare positive integer. Hence, applying (i) of Lemma 2.1 to (3.10), there exists a positive integer n' which makes

$$b^{y/2} + g\sqrt{ad} = \left(u_1 + v_1\sqrt{ad}\right)^{n'}, \quad (3.11)$$

where (u_1, v_1) is the least solution of (3.9).

For any positive integer n , let

$$u_n + v_n\sqrt{ad} = \left(u_1 + v_1\sqrt{ad}\right)^n. \quad (3.12)$$

If $2 \mid n'$, then from (3.11) and (3.12) we get $b^{y/2} = u_{n'}$ and, by (ii) of Lemma 2.1, $b \equiv \pm 1 \pmod{8}$, which contradicts the assumption. So we get

$$2 \nmid n'. \quad (3.13)$$

Since $2 \nmid x$, we see from the first equality of (3.2) that the equation

$$aU^2 - dV^2 = 1, \quad U, V \in \mathbb{N} \quad (3.14)$$

has a solution

$$(U, V) = \left(a^{(x-1)/2}, f\right). \quad (3.15)$$

Let (U_1, V_1) be the least solution of (3.14). For any odd positive integer m , let

$$U_m\sqrt{a} + V_m\sqrt{d} = \left(U_1\sqrt{a} + V_1\sqrt{d}\right)^m. \quad (3.16)$$

Applying (i) of Lemma 2.2 to (3.15), by (3.16), there exists an odd positive integer m' which makes

$$\left(a^{(x-1)/2}, f\right) = (U_{m'}, V_{m'}). \quad (3.17)$$

Hence, by Lemma 2.3, we get from (3.1) and (3.17) that

$$\text{ord}_2(aU_1^2) = \text{ord}_2(aU_{m'}^2) = \text{ord}_2(a^x) \geq x \geq 3. \quad (3.18)$$

By (ii) of Lemma 2.2, we find from (3.11), (3.13) and (3.16) that

$$\begin{aligned} b^{y/2} + g\sqrt{ad} &= \left(U_1\sqrt{a} + V_1\sqrt{d}\right)^{2n'} = \left(\left(U_1\sqrt{a} + V_1\sqrt{d}\right)^{n'}\right)^2 \\ &= \left(U_{n'}\sqrt{a} + V_{n'}\sqrt{d}\right)^2. \end{aligned} \quad (3.19)$$

Since $aU_{n'}^2 - dV_{n'}^2 = 1$, by (3.19), we have

$$b^{y/2} = aU_{n'}^2 + dV_{n'}^2 = 2aU_{n'}^2 - 1. \quad (3.20)$$

Further, by Lemma 2.3, we have $\text{ord}_2(aU_{n'}^2) = \text{ord}_2(aU_1^2)$. Hence, by (3.18), we get $\text{ord}_2(aU_{n'}^2) \geq 3$ and $aU_{n'}^2 \equiv 0 \pmod{8}$. Therefore, by (3.20), we obtain $b^{y/2} \equiv 7 \pmod{8}$. But, since $b \equiv \pm 3 \pmod{8}$, it is impossible. Thus, the theorem is proved.

4. Proof of Theorem 1.2

In this section, we assume that $a = 2$, $b \equiv \pm 3 \pmod{8}$ and (x, y, z) is a solution of (1.4) with $2 \nmid x$. By Theorem 1.1, we have

$$x = 1. \quad (4.1)$$

Since $a = 2$, substituting (4.1) into (3.2), we get

$$d = f = 1 \quad (4.2)$$

and

$$b^y - 1 = 2g^2, \quad z = g, \quad g \in \mathbb{N}. \quad (4.3)$$

If $b \equiv 5 \pmod{8}$, then from the first equality of (4.3) we get $1 = (-2/b) = (2/b) = -1$, a contradiction, where $(*/b)$ is the Jacobi symbol. Therefore, if $a = 2$ and $b \equiv 5 \pmod{8}$, then (1.4) has no solutions (x, y, z) with $2 \nmid x$.

We just need to consider the case $b \equiv 3 \pmod{8}$. Applying Lemma 2.9 to the first equality of (4.3), by (4.1) and (4.3), equation (1.4) has only the solution

$$b = 3, \quad (x, y, z) = (1, 5, 11) \quad (4.4)$$

with $y > 2$.

When $y = 2$, by the first equality of (4.3), $(u, v) = (b, g)$ is a solution of (2.5). Since $(u_1, v_1) = (3, 2)$ is the least solution of (2.5), by (i) of Lemma 2.1, we get from (1.5) that

$$(b, g) = (r_{n'}, s_{n'}), \quad n' \in \mathbb{N}. \quad (4.5)$$

Further, since $b \equiv 3 \pmod{8}$, by Lemma 2.4, we see from (4.5) that $2 \nmid n'$. Hence, by (4.1), (4.2), (4.3) and (4.5), we obtain

$$b = r_m, \quad (x, y, z) = (1, 2, s_m), \quad m \in \mathbb{N}, \quad 2 \nmid m. \quad (4.6)$$

When $y = 1$, by (4.1), (4.2) and (4.3), we have

$$b = 2g^2 + 1, \quad (x, y, z) = (1, 1, g), \quad g \in \mathbb{N}, \quad 2 \nmid g. \quad (4.7)$$

Thus, since $r_1 = 2 \cdot 1^2 + 1 = 3$, the combination of (4.4), (4.6) and (4.7) yields the solutions (i), (ii) and (iii). The theorem is proved.

5. Proof of Theorem 1.3

By Theorem 1.2, we get $N(2, 3) = 3$ immediately. By Lemma 2.5, if $b = 2g^2 + 1$ and $g = R_m$ with $m > 1$, then $b = r_m > 3$. Hence, by Theorem 1.2, we have $N(2, b) = 2$. In addition, if $b = 2g^2 + 1$ with $g \neq R_m$ or $b \neq 2g^2 + 1$, then $N(2, b) = 1$ or 0 . The theorem is proved.

6. Proof of Theorems 1.4 and 1.5

Proof of Theorem 1.4. By Theorem 1.1, we may assume that $x = 2x_0$ for some $x_0 \in \mathbb{N}$. In addition, since a is a square, we may write $a = a_0^2$ for some $a_0 \in \mathbb{N}$. Then, by the first equality of (3.2), we get

$$(a_0^{x_0})^4 - df^2 = 1. \quad (6.1)$$

It is clear from (6.1) that

$$d \text{ is not a square.} \quad (6.2)$$

Applying Lemma 2.8 to (6.1), we see that either $a^{x_0} = u'_1$ or $a^{x_0} = 2(u'_1)^2 - 1$, where (u'_1, v'_1) is the least solution of (2.1) with $D = d$. Since $2 \mid a$, we must have

$$a^{x_0} = u'_1. \quad (6.3)$$

On the other hand, we know by $4 \mid a$ and $2 \mid x$ that (3.4) holds, which together with (3.6) and (3.7) yields $2 \mid y$. Since $a = a_0^2$, we see from the second equality of (3.2) that (2.1) with $D = d$ has a solution $(u, v) = (b^{y/2}, a_0g)$. By (i) of Lemma 2.1 and (6.2), we have

$$(u'_n, v'_n) = (b^{y/2}, a_0g), \quad n \in \mathbb{N}, \quad (6.4)$$

where $u'_n + v'_n\sqrt{d} = (u'_1 + v'_1\sqrt{d})^n$. If $2 \mid n$, then, by (ii) of Lemma 2.1, $b \equiv \pm 1 \pmod{8}$, which contradicts the assumption. If $2 \nmid n$, then, by (iii) of Lemma 2.1, $u'_1 \mid u'_n$. However, by (6.3) and (6.4), we have $a \mid b^{y/2}$, which contradicts $2 \mid a$ and $b \equiv \pm 3 \pmod{8}$. The theorem is proved. \square

Proof of Theorem 1.5. By Theorem 1.4, we have

$$x = 1. \quad (6.5)$$

(i) Substituting $a = 4$ and (6.5) into (3.2), we get

$$d = 3, \quad f = 1$$

and

$$b^y - 1 = 12g^2, \quad z = 3g, \quad g \in \mathbb{N}. \quad (6.6)$$

Obviously, we have $b \neq 3$. If b has a prime divisor p with $p \equiv 11 \pmod{24}$, then by (6.6) we have

$$-1 = \left(\frac{-1}{p}\right) = \left(\frac{12g^2}{p}\right) = \left(\frac{3}{p}\right) = 1,$$

a contradiction. Thus, (i) is proved.

(ii) Substituting $a = 16$ and (6.5) into (3.2), we get

$$d = 15, \quad f = 1$$

and

$$b^y - 1 = 15 \cdot 16g^2, \quad z = 15g, \quad g \in \mathbb{N}. \quad (6.7)$$

Obviously, we have $b \notin \{3, 5\}$. If b has a prime divisor p with $p \equiv 11, 43, 59$ or $67 \pmod{120}$, then, by (6.7),

$$-1 = \left(\frac{-1}{p}\right) = \left(\frac{15}{p}\right) = 1,$$

a contradiction. If b has a prime divisor p with $p \equiv 13, 29, 37$ or $101 \pmod{120}$, then, by (6.7),

$$1 = \left(\frac{-1}{p}\right) = \left(\frac{15}{p}\right) = -1,$$

a contradiction. Thus, the theorem is proved. \square

Acknowledgements. The authors thank the referee for careful reading and helpful comments.

References

- [1] J. H. E. COHN: *The Diophantine equation $(a^n - 1)(b^n - 1) = x^2$* , Period. Math. Hung. 44 (2002), pp. 169–175, DOI: [10.1023/A:1019688312555](https://doi.org/10.1023/A:1019688312555).
- [2] M.-H. LE: *A necessary and sufficient condition for the equation $x^4 - Dy^2 = 1$ to have positive integer solutions*, Chinese Sci. Bull. 30 (1984), p. 1698.
- [3] M.-H. LE, G. SOYDAN: *A brief survey on the generalized Lebesgue-Ramanujan-Nagell equation*, Surv. Math. Appl. 15 (2020), pp. 473–523.
- [4] L. LI, L. SZALAY: *On the exponential Diophantine equation $(a^n - 1)(b^n - 1) = x^2$* , Publ. Math. Debrecen 77 (2010), pp. 465–470, DOI: [10.5486/PMD.2010.4697](https://doi.org/10.5486/PMD.2010.4697).
- [5] L. J. MORDELL: *Diophantine equations*, London: Academic Press, 1969.
- [6] T. NAGELL: *Sur l'impossibilité de quelques équations à deux indéterminées*, Norsk Mat. Forenings Skr. 13 (1923), pp. 65–82.
- [7] L. SZALAY: *On the Diophantine equation $(2^n - 1)(3^n - 1) = x^2$* , Publ. Math. Debrecen 57 (2000), pp. 1–9, DOI: [10.5486/PMD.2000.2069](https://doi.org/10.5486/PMD.2000.2069).
- [8] R.-Z. TONG: *On the Diophantine equation $(2^x - 1)(p^y - 1) = 2z^2$* , Czech. Math. J. 71 (2021), pp. 689–696, DOI: [10.21136/CMJ.2021.0057-20](https://doi.org/10.21136/CMJ.2021.0057-20).
- [9] R. W. VAN DER WAALL: *On the Diophantine equations $x^2 + x + 1 = 3y^2$, $x^3 - 1 = 2y^2$, $x^3 + 1 = 2y^2$* , Simon Stevin 46 (1972/1973), pp. 39–51.
- [10] D. T. WALKER: *On the Diophantine equation $mx^2 - ny^2 = \pm 1$* , Amer. Math. Monthly 74 (1967), pp. 504–513, DOI: [10.1080/00029890.1967.11999992](https://doi.org/10.1080/00029890.1967.11999992).